

# First-Year Electromagnetism: Problem Set 2

Hilary Term 2015, Prof LM Herz

## B. The Method of Image Charges

**B.0 Background.** Explain the factors that determine the parallel and normal components of the electric field near the surface of a flat metal plate.

**B.1 Charge monopole near a flat metal surface.** A point charge  $q$  is placed at a perpendicular distance  $d$  from a point  $O$  on a flat, infinite plate that is conducting and earthed.

- Sketch the resulting lines of the electric field and calculate the force  $F$  between the charge and the plate.
- Use the method of images to show that the magnitude  $E$  of the electric field at the point  $P$ , a distance  $r$  along the plane from  $O$ , is

$$E(r) = \frac{qd}{2\pi\epsilon_0(r^2 + d^2)^{3/2}}.$$

- Show that the charge on the plane is  $-q$ .
- Find the work done in moving the charge to an infinite distance from the plane. Hence find the minimum energy an electron must have in order to escape from a metal surface (assume that it starts at a distance 0.1 nm, which is about one atomic diameter above the surface). Express your answer in electron volts.

**B.2 Two charges near a flat metal surface.** Two charges  $+Q$  and  $-Q$  are a horizontal distance  $a$  apart and a vertical distance  $b$  above a large conducting sheet. Find the components of the forces acting on each charge.

**B.3 Charge monopole near two orthogonal metal surfaces.** Two semi-infinite plane conducting plates are joined together at right angle. A charge  $Q$  is situated near the join at a distance  $a$  from each plate.

- Show that the electric field is zero at any point along the join.
- Find the field just above the surface of one of the plates at the point closest to the charge.
- Sketch the equipotentials near the charge and near the plate.
- Calculate the surface charge density on the metal plates at the points closest to the charge  $Q$ .

(Hint: you need to consider three image charges.)

**B.4 Uniformly charged rod near a metal surface.** An infinite, thin, uniformly charged rod (line charge density  $\lambda$ ) is placed parallel to a metal plate a distance  $d$  above it. Use the result obtained in Problem A.4 to calculate the electric field at a point  $P$  close to the surface of the plate as a function of the distance  $\overline{PM}$ , where  $M$  is the closest midpoint between the charge and the image charge.

## C. Electric Fields derived from Gauss' Law

**C.0 Background.** State Gauss' Law and explain how it may be used to determine the electric field arising from a spherically symmetric charge density distribution  $\rho(r)$ .

### C.1 Uniformly charged sphere.

(a) Charge  $+q$  is distributed uniformly throughout the volume of a sphere of radius  $a$ . Show that the electric field  $\mathbf{E}$  and potential  $V$  at a distance  $r$  from the centre of the sphere are given by:

$$\mathbf{E} = \begin{cases} \frac{qr}{4\pi\epsilon_0 a^3} \hat{\mathbf{r}} & \text{and} \\ \frac{q}{4\pi\epsilon_0 r^2} \hat{\mathbf{r}} & \end{cases} \quad V = \begin{cases} \frac{q}{4\pi\epsilon_0 a^3} \left( \frac{3a^2}{2} - \frac{r^2}{2} \right) & \text{for } 0 \leq r \leq a \\ \frac{q}{4\pi\epsilon_0 r} & \text{for } a \leq r \end{cases}$$

(b) Repeat the calculation of fields and potentials for the charge now being uniformly distributed over the surface of a sphere of radius  $a$ .  
 (c) Draw graphs of the electric fields (magnitude) and the potentials for both cases (solid sphere and shell). Take care to illustrate the relation  $E = -(\partial V / \partial r)$  everywhere and account for any discontinuities that occur at  $r = a$ .

**C.2 Coulomb energy of the nucleus.** The nucleus of an atom can be considered to be a charge  $+Ze$  uniformly distributed throughout a sphere of radius  $a$ .

(a) Show that the potential energy  $W$  of a nucleus arising from the assembly of its charge is given by  $W = 3(Ze)^2(20\pi\epsilon_0 a)^{-1}$ .  
 (b) What would the potential energy be if the charge was instead spread uniformly over the surface of the nucleus?

**C.3 Electron in a hydrogen atom.** From a quantum mechanical treatment, the potential experienced by an electron in a hydrogen atom at a distance  $r$  from the nucleus is

$$V = \frac{q}{4\pi\epsilon_0} \left( \frac{\exp(-2r/a) - 1}{r} + \frac{\exp(-2r/a)}{a} \right)$$

where  $a$  is a constant and is a measure of the “size” of the atom.

(a) Sketch  $V(r)$  for  $0 \leq r \leq \infty$  and comment on the shape of the curve.  
 (b) Find the magnitude of the electric field at a distance  $r \ll a$  from the nucleus.  
 (c) Show that, when an external electric field  $E_{ext}$  is applied, the atom develops a dipole moment of magnitude  $p$  (you may assume that the electron cloud remains spherical and merely moves relative to the nucleus). By considering the force on the nucleus, calculate  $p$  and show that the polarisability  $p/E_{ext}$  is equal to  $3\pi\epsilon_0 a^3$ .  
 (d) For the hydrogen atom,  $a = 0.5 \times 10^{-10}$  m. Show that even for the largest accessible fields of  $\sim 10^6$  V m $^{-1}$  the electron charge cloud moves relative to the nucleus by only about  $10^{-17}$  m (which justifies the assumption  $r \ll a$ ).  
 (e) Use Gauss law to calculate the total charge in the cloud.